

A Supersymmetric Contribution to the Neutrino Mass Matrix and Breaking of μ - τ Symmetry

Naoyuki Haba^{a,*} and Werner Rodejohann^{b,†}

^a *University of Tokushima,
Institute of Theoretical Physics,
Tokushima-shi 770-8502, Japan*

^b *Physik-Department, Technische Universität München,
James-Frank-Straße,
85748 Garching bei München, Germany*

Supersymmetry broken by anomaly mediation suffers from tachyonic slepton masses. A possible solution to this problem results in “decoupling”, i.e., the first two generations of sfermions are much heavier than the third one. We note that in this scenario a sizable loop-induced contribution to the neutrino mass matrix results. As an application of this scenario we take advantage of the fact that the decoupling evidently not obeys 2–3 generation exchange symmetry. In the neutrino sector, this 2–3 symmetry (or μ - τ symmetry) is a useful Ansatz to generate zero θ_{13} and maximal θ_{23} . The induced deviations from these values are given for some examples, thereby linking SUSY breaking to the small parameters (including possibly the solar mass splitting) of the neutrino sector.

PACS numbers: 11.30.Pb, 14.60.Pq

Keywords: SUSY breaking; neutrino mixing; μ - τ symmetry

Supersymmetry (SUSY) is the best motivated candidate for beyond the Standard Model (SM) physics, since it for instance provides Dark Matter candidates or solves the gauge hierarchy problem. Phenomenologically, the requirement of SUSY breaking is obvious, its origin and mechanism are however still a mystery. One possibility is the anomaly mediated SUSY breaking (AMSB) [1], which always exists in supergravity frameworks. It avoids the presence of sizable flavor changing neutral currents, the so-called SUSY flavor problem, but has a well-known drawback, namely the presence of tachyonic slepton masses. One simple solution for this dilemma is obtained by introducing an additional D -term contribution to the sfermion masses [2]. Taking into account this contribution can have an interesting effect: the 1st and 2nd generation sfermion masses are heavy, $\mathcal{O}(1 \text{ TeV})$, while the 3rd generation sfermion and gaugino masses are $\mathcal{O}(100 \text{ GeV})$. This latter scale corresponds to the scale of the soft SUSY breaking masses induced by AMSB, $\simeq 0.01 m_{3/2}$ (with $m_{3/2}$ the gravitino mass). We denote this framework here the “AMSB-decoupling” scenario and will place our analysis there. In the AMSB scenario the 1-loop correction to dimension 5 operators can be of the same order than the tree level contribution [3]. One example is the dimension five operator responsible for neutrino masses (see Fig. 1):

$$\mathcal{L}_{m_\nu} = \frac{\kappa_{ij}}{2M_R} (\psi_{L_i} H) (\psi_{L_j} H) + \text{h.c.} \quad (1)$$

Here ψ_{L_i} is the lepton doublet of flavor i , H the up-type Higgs doublet and M_R the scale of neutrino mass generation. After electroweak symmetry breaking this operator is the neutrino mass matrix $m_\nu = \kappa \langle H \rangle^2 / M_R$. In the present work we will show explicitly how in the AMSB-decoupling framework a sizable and flavor-dependent contribution to Eq. (1) arises and will also discuss some interesting applications.

The structure of κ defines the neutrino mass and mixing phenomena [4] and explaining their peculiar scheme is one of the most interesting problems of particle physics. An interesting Ansatz is to implement a 2–3 or μ - τ exchange symmetry [5] in κ . This yields zero θ_{13} and maximal θ_{23} , which are the best-fit points of global analyses of neutrino data [6]. Deviations from these values are achieved by breaking of μ - τ symmetry, which however is usually done by hand [7, 8]. We propose here to take advantage of the AMSB-decoupling contribution to a neutrino mass matrix with μ - τ symmetry. The fact that the first two sfermion families are much heavier than the third one apparently violates 2–3 exchange symmetry. Assuming that κ conserves μ - τ symmetry and taking the μ - τ violating AMSB correction to κ into account leads to testable corrections to the initial mixing parameters. The size of the contribution to Eq. (1) depends on the SUSY masses. Hence, the implied small values of θ_{13} and $\theta_{23} - \pi/4$ are also sensitive to the SUSY masses. In addition, it is also possible to generate the small solar neutrino mass squared difference via the AMSB-decoupling term.

Now let us estimate the 1-loop correction for the neutrino dimension 5 operator in Eq. (1). As an effect of

*Electronic address: haba@ph.tum.de

†Electronic address: werner.rodejohann@ph.tum.de




FIG. 1: The dimension 5 operator responsible for neutrino masses (left, Eq. (1)) and the contribution in the AMSB-decoupling scenario (right, Eq. (3)).

quantum corrections, the neutrino mass term is obtained by replacing every entry ij as follows:

$$(m_\nu)_{ij} \rightarrow (m_\nu)_{ij} (1 + \Delta_{ij}) \text{ for all } i, j. \quad (2)$$

Here the factor Δ_{ij} is the correction from the (neutral) wino and bino dressed 1-loop diagram (see Fig. 1),

$$\begin{aligned} \Delta_{ij} &\simeq \frac{1}{2} \left(\frac{\alpha_2}{4\pi} \right) F_\phi M_2^* f_{ij}^{(2)} + \frac{3}{10} \left(\frac{\alpha_1}{4\pi} \right) F_\phi M_1^* f_{ij}^{(1)} \\ &\simeq \frac{|M_2|^2}{2} (f_{ij}^{(1)} + f_{ij}^{(2)}). \end{aligned} \quad (3)$$

We defined the loop-function $f_{ij}^{(I)} = f(M_I, \tilde{m}_{L_i}, \tilde{m}_{L_j})$ as

$$f(M_I, a, b) = \frac{1}{a^2 - b^2} \left[\frac{a^2}{a^2 - |M_I|^2} \ln \frac{a^2}{|M_I|^2} - (a \leftrightarrow b) \right], \quad (4)$$

where $\alpha_{1,2}$ are the gauge couplings, $M_{1,2}$ the wino ($\tilde{W}_3$) and bino ($\tilde{b}$) masses, and $\tilde{m}_{L_i}$ denotes the mass of the i th generation slepton. F_ϕ is the F -component of the compensating multiplet of the AMSB sector and is of the order of the gravitino mass. The second equation in Eq. (3) uses the AMSB relation $M_I = (\alpha_I/4\pi) b_I F_\phi$ ($I = 1, 2$ and $b_1 = 33/5$ and $b_2 = 1$ are the beta functions), which implies $M_2/M_1 \simeq 0.3$. In the decoupling mass spectrum under consideration, the 1-3 and 2-3 generation sfermion mixings are small, so that we can safely neglect their mixings at the vertices. Note that we did not specify the origin of the dimension 5 operator κ . Our analysis is therefore independent of the leading mechanism of neutrino mass generation.

The contribution to Δ_{ij} is of order $\alpha_I/(4\pi) M_I F_\phi f_{ij}^I$, with $f_{ij}^I \sim M^{-2}$, where M is here some soft SUSY breaking mass. In other SUSY breaking scenarios there are also diagrams as the one displayed in Fig. 1. However, such scenarios typically predict that F_ϕ , the gravitino, gaugino and sfermion masses are of the same order of magnitude. As a consequence, the loop-diagram gives a contribution to κ strongly suppressed by a loop factor of 10^{-2} . The crucial point we wish to make here occurs

when there is *only* the AMSB breaking effect in which case F_ϕ is larger by a factor of 10^2 than the gaugino masses. Then this factor 10^2 cancels with $\alpha_I/(4\pi)$ and the contribution of Δ_{ij} to κ_{ij} is sizable.

Let us show two typical examples for the magnitude of Δ_{ij} . When we take $M_2 = 220$ GeV, $\tilde{m}_{L_3} = 300$ GeV, and $\tilde{m}_{L_{1,2}} \geq 4.5$ TeV (“small slepton masses”), the dimensionless matrix Δ_{ij} is estimated as

$$\Delta_{ij} \simeq \begin{pmatrix} \lambda^4 & \lambda^4 & \lambda^3 \\ \cdot & \lambda^4 & \lambda^3 \\ \cdot & \cdot & \lambda \end{pmatrix} \simeq \begin{pmatrix} 0 & 0 & 0 \\ \cdot & 0 & 0 \\ \cdot & \cdot & \lambda \end{pmatrix}. \quad (5)$$

We introduced here the small parameter $\lambda = 0.22$, which is of the order of the Cabibbo angle, and proves very useful in the estimates. The heavier the 1st and 2nd generation slepton masses become, the smaller the elements except for the (3, 3) entry become. It is the peculiar mass ordering of the sfermions that makes the (3, 3) entry dominate in Δ_{ij} .

A similar hierarchy in Δ_{ij} , but with a smaller (3, 3) entry is obtained by choosing $M_2 = 200$ GeV, $\tilde{m}_{L_3} = 760$ GeV, and $\tilde{m}_{L_{1,2}} \geq 8.8$ TeV (“large slepton masses”), for which

$$\Delta_{ij} \simeq \begin{pmatrix} \lambda^5 & \lambda^5 & \lambda^4 \\ \cdot & \lambda^5 & \lambda^4 \\ \cdot & \cdot & \lambda^2 \end{pmatrix} \simeq \begin{pmatrix} 0 & 0 & 0 \\ \cdot & 0 & 0 \\ \cdot & \cdot & \lambda^2 \end{pmatrix}. \quad (6)$$

We can therefore adjust the magnitude of the correction to the neutrino mass matrix by choosing the slepton and gaugino mass spectrum. The sizable differences of the slepton masses and the light gauginos (as typical for AMSB scenarios) might be demonstrated at future colliders such as LHC or the ILC.

As obvious from Eqs. (5, 6), the correction to m_ν violates μ - τ symmetry and can serve as a perturbation to such mixing scenarios. We will illustrate this in what follows. A general μ - τ symmetric neutrino mass matrix leaves the neutrino mass spectrum and the solar neutrino mixing angle unconstrained. To be definite, we assume here first a normal hierarchy and bimaximal mixing [9], for which $\theta_{12} = \pi/4$. The mass matrix (with the smallest neutrino mass $m_1 \simeq 0$) can be written as

$$m_\nu \simeq \frac{m_3}{2} \begin{pmatrix} \epsilon & \epsilon/\sqrt{2} & \epsilon/\sqrt{2} \\ \cdot & 1 + \epsilon/2 & -1 + \epsilon/2 \\ \cdot & \cdot & 1 + \epsilon/2 \end{pmatrix}, \quad (7)$$

with $\epsilon \simeq \sqrt{\Delta m_\odot^2 / \Delta m_A^2}$, where the square root lies at 3 (1) σ between 0.15 and 0.25 (0.17 and 0.21), with a best-fit value of 0.19 [6]. The effect of the AMSB-decoupling contribution in Eq. (5) is now basically just multiplying the (3, 3) entry of Eq. (7) with $(1 + \lambda)$. We can then

predict the modified neutrino observables:

$$R \equiv \frac{\Delta m_{\odot}^2}{\Delta m_A^2} \simeq \frac{4\epsilon^2 + \epsilon\lambda + \lambda^2/8}{4 + 2\lambda} \quad , \quad |U_{e3}| \simeq \frac{\epsilon\lambda}{8} \quad (8)$$

$$\tan \theta_{23} \simeq 1 - \lambda/2 \quad , \quad \tan 2\theta_{12} \simeq \frac{4\epsilon}{\lambda} \quad ,$$

which for $\lambda = 0$ reproduces bimaximal mixing with $\Delta m_{\odot}^2 = \epsilon^2 \Delta m_A^2$. For $\lambda \lesssim \epsilon \simeq \sqrt{R}$ the required large solar neutrino mixing (observation indicates $\tan 2\theta_{12} \simeq 2\sqrt{2}$) is easily achieved. Note that the deviation from maximal atmospheric mixing is large, namely of order $\sqrt{R}$, whereas $|U_{e3}|$ is of order R . Both quantities depend on the SUSY breaking parameters as encoded in λ .

Another possibility for bimaximal mixing and the normal hierarchy is when before breaking one chooses $m_1 = -m_2$, i.e., we start with vanishing solar Δm^2 . The mass matrix reads

$$m_{\nu} \simeq \frac{m_3}{2} \begin{pmatrix} 0 & \sqrt{2}\epsilon & \sqrt{2}\epsilon \\ \cdot & 1 & -1 \\ \cdot & \cdot & 1 \end{pmatrix} \quad , \quad (9)$$

with $\epsilon = m_2/m_3 \ll 1$. Note that here the effective mass as measurable in neutrino-less double beta decay (the (1,1) element of m_{ν}) is much smaller than in the previous example. Taking now the perturbation from Eq. (5) into account yields

$$R \simeq \frac{2\epsilon\lambda}{4 + 2\lambda} \quad , \quad |U_{e3}| \simeq \frac{\epsilon\lambda}{4} \quad , \quad \tan 2\theta_{12} \simeq \frac{8\epsilon}{\lambda}$$

and an identical results for θ_{23} as in Eq. (8). We stress that in this scenario there is a link between the small solar mass splitting and the breaking of supersymmetry. We can also perturb tri-bimaximal mixing [10], which is μ - τ symmetry with $\sin^2 \theta_{12} = \frac{1}{3}$. This requires a smaller perturbation to the mass matrix, i.e., Eq. (6), and therefore larger slepton masses. Initial tri-bimaximal mixing with $m_1 = 0$ corrected by Eq. (6) yields

$$\sin^2 \theta_{12} \simeq \frac{1}{3} + \frac{1}{9} \lambda^2 / \epsilon \quad , \quad (10)$$

very small $|U_{e3}| \simeq \epsilon \lambda^2 / \sqrt{72}$ and moreover close-to-maximal $\tan \theta_{23} \simeq 1 - \lambda^2/2$. R is of order ϵ^2 . All deviations from the initial values are much smaller than for initial bimaximal mixing. In principle it would be possible that the correction to the mass matrix stems from Eq. (5), i.e., is larger. To accommodate the observed value of $\sin^2 \theta_{12} \simeq \frac{1}{3}$, however, the contribution proportional to λ should be suppressed by fine-tuned values of possible CP phases.

Consider now the inverted hierarchy. Since in the inverted hierarchy one can not perturb the bimaximal mixing to accommodate the observed large but non-maximal solar mixing (unless one accepts fine-tuning), we start

with free θ_{12} :

$$m_{\nu} \simeq m \begin{pmatrix} A & B & B \\ \cdot & D & D \\ \cdot & \cdot & D \end{pmatrix}$$

$$\text{with } A \equiv s_{12}^2 - (1 - \epsilon) c_{12}^2 \quad , \quad B \equiv \frac{1}{2\sqrt{2}} (2 - \epsilon) \sin 2\theta_{12} \quad ,$$

$$D \equiv \frac{\epsilon}{4} + \frac{1}{4} (2 - \epsilon) \cos 2\theta_{12}$$

and $\epsilon \equiv (m_2 - m_1)/m$, where $m_2 \simeq m_1 \simeq m \simeq \sqrt{\Delta m_A^2}$. The two non-zero masses m_1 and m_2 have opposite CP parities here. As in the normal hierarchy with tri-bimaximal mixing, we require a small perturbation to the mass matrix: if we would add Eq. (5) to this matrix, then the ratio of solar and atmospheric Δm^2 would be of order $\lambda + 2\epsilon$, which is too large (this will also hold for the unstable case of equal CP parities of the masses m_1 and m_2). Hence, we are lead to use Eq. (6), corresponding to large slepton masses. Choosing θ_{12} such that $\sin^2 \theta_{12} = \frac{1}{3}$, i.e., again tri-bimaximal mixing [10], one finds

$$R \simeq \frac{\lambda^2 + 2\epsilon}{1 + \frac{2}{3}\lambda^2 + \frac{8}{9}\epsilon} \quad , \quad \tan^2 \theta_{23} \simeq 1 + \frac{2}{3}\lambda^2 \quad ,$$

$$|U_{e3}| \simeq \frac{\sqrt{2}\lambda^2}{3} \quad , \quad \sin^2 \theta_{12} \simeq \frac{1}{3} - \frac{1}{9}\lambda^2 + \frac{2}{27}\epsilon \quad . \quad (11)$$

The order of magnitude of θ_{13} is the same as in the case of normal hierarchy and initial bimaximal mixing, namely of order R . The deviation from maximal atmospheric mixing is of order R . Starting with $\epsilon = 0$, i.e., with vanishing $\Delta m_{\odot}^2$, we have $R \simeq 3|U_{e3}|/\sqrt{2}$ and

$$\sin^2 \theta_{12} \simeq \frac{1}{3} - \frac{1}{3\sqrt{2}} |U_{e3}| \simeq \frac{1}{2} - \frac{1}{6} \tan^2 \theta_{23} \quad . \quad (12)$$

Again, a large contribution to the mass matrix corresponding to smaller slepton masses is in principle possible but requires fine-tuned CP phases.

Up to now it is obvious that the μ - τ symmetry was only obeyed by the neutrino mass matrix, while the charged lepton mass matrix was real and diagonal. We choose now an example in which both the neutrinos and the charged leptons conserve μ - τ symmetry. Naively, one would expect that even after breaking the symmetry the PMNS matrix would contain only a sizable 12 mixing, since the (close-to-)maximal 23 angle cancels in the product of the matrices diagonalizing the neutrinos and charged leptons, respectively. This can change for quasi-degenerate light neutrinos [8]. The mass matrices are:

$$m_{\ell,\nu} = \begin{pmatrix} A_{\ell,\nu} & B_{\ell,\nu} & B_{\ell,\nu} \\ \cdot & D_{\ell,\nu} + E_{\ell,\nu} & E_{\ell,\nu} - D_{\ell,\nu} \\ \cdot & \cdot & D_{\ell,\nu} + E_{\ell,\nu} \end{pmatrix},$$

$$\text{with } A_{\ell,\nu} \equiv m_{e,1} (c_{12}^{\ell,\nu})^2 + m_{\mu,2} (s_{12}^{\ell,\nu})^2,$$

$$B_{\ell,\nu} \equiv (m_{\mu,2} - m_{e,1}) c_{12}^{\ell,\nu} s_{12}^{\ell,\nu} / \sqrt{2},$$

$$D_{\ell,\nu} \equiv m_{\tau,3}/2, \quad E_{\ell,\nu} \equiv \frac{1}{2}(m_{\mu,2} (c_{12}^{\ell,\nu})^2 + m_{e,1} (s_{12}^{\ell,\nu})^2).$$

Here $m_{e,\mu,\tau}$ are the charged lepton masses and $c_{12}^{\ell,\nu}$ ($s_{12}^{\ell,\nu}$) the cosine (sine) of the 12 mixing angle which diagonalizes the charged lepton and neutrino mass matrix, respectively. Note that m_ℓ is symmetric. That does not affect our analysis, since for non-symmetric m_ℓ one would have to choose a μ - τ symmetric $m_\ell m_\ell^\dagger$, which would give the same results. Perturbing the neutrino mass matrix with the term of Eq. (5), we have in the basis of diagonal charged leptons (choosing for simplicity $\theta_{12}^\ell \ll 1$)

$$m_\nu \simeq \begin{pmatrix} A_\nu & \sqrt{2} B_\nu & 0 \\ \cdot & 2 E_\nu & \frac{1}{2}(D_\nu + E_\nu) \lambda \\ \cdot & \cdot & 2 D_\nu \end{pmatrix}, \quad (13)$$

where only the leading terms are shown. Large atmospheric neutrino mixing is only possible for $D_\nu \simeq E_\nu$. This corresponds to a quasi-degenerate spectrum and is exactly the scenario put forward in [8] to accommodate μ - τ symmetry in both the charged lepton and the neutrino mass matrix. Focusing on the 23 sector and denoting the common mass scale with m_0 , close-to-maximal θ_{23} is achieved for $\lambda (D_\nu + E_\nu) \gg |D_\nu - E_\nu|$, which leads to $\cos 2\theta_{23} \simeq 4 m_0 |D_\nu - E_\nu| / \Delta m_A^2$ [13]. The parameter λ is related to the masses:

$$\lambda \simeq \frac{\Delta m_A^2}{2m_0^2} \simeq \begin{cases} 0.11 & \text{for } m_0 = 0.1 \text{ eV}, \\ 0.012 & \text{for } m_0 = 0.3 \text{ eV}. \end{cases} \quad (14)$$

Such values of the neutrino masses are in agreement even with the tightest cosmological constraints [11]. Light neutrino masses around 0.1 eV require therefore a perturbation with small slepton masses and larger neutrino masses need larger slepton masses.

Numerically, for $m_0 \simeq 0.1$ eV we have $\cos 2\theta_{23} \simeq |D_\nu - E_\nu| / (0.0055 \text{ eV})$, which shows the required fine-tuning in such a scenario. If $\theta_{23} = \pi/4 - 0.1$, then $|D_\nu - E_\nu| \simeq 0.0011$ eV and for $\theta_{23} = \pi/4 - 0.01$ one has $|D_\nu - E_\nu| \simeq 0.0001$ eV. For a larger mass of $m_0 \simeq 0.3$ eV one finds that for $\theta_{23} = \pi/4 - 0.1$ one requires $|D_\nu - E_\nu| \simeq 0.0004$ eV, while for $\theta_{23} = \pi/4 - 0.01$ one has $|D_\nu - E_\nu| \simeq 0.00004$ eV.

In summary, anomaly mediated supersymmetry breaking with connection to decoupling of the sfermions can generate a sizable contribution to the neutrino mass matrix. This can serve as a natural explanation for a perturbation of neutrino mixing scenarios. Noting that the decoupling scenario violates 2-3 exchange symmetry, it is natural to take μ - τ symmetry as an example. We showed in various examples that interesting and testable [12] correlations between the neutrino mixing parameters result. We stress again that in the scenarios presented here the breaking of supersymmetry is intimately related to small U_{e3} , $\cos 2\theta_{23}$ and even to the small ratio of the solar and atmospheric Δm^2 .

Acknowledgments

The work of N.H. was supported by the Alexander-von-Humboldt-Foundation. The work of W.R. was supported by the ‘‘Deutsche Forschungsgemeinschaft’’ in the ‘‘Sonderforschungsbereich 375 f ur Astroteilchenphysik’’ and under project number RO-2516/3-1. We would like to thank T. Ota and N. Okada for useful discussions.

-
- [1] L. Randall and R. Sundrum, Nucl. Phys. B **557**, 79 (1999); G. F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, JHEP **9812**, 027 (1998).
 - [2] I. Jack and D. R. Jones, Phys. Lett. B **482**, 167 (2000); N. Kitazawa, N. Maru and N. Okada, Phys. Rev. D **63**, 015005 (2001); N. Arkani-Hamed, D. E. Kaplan, H. Murayama and Y. Nomura, JHEP **0102**, 041 (2001).
 - [3] N. Haba and N. Okada, hep-ph/0601003.
 - [4] R. N. Mohapatra *et al.*, hep-ph/0510213.
 - [5] For instance, C. S. Lam, Phys. Lett. B **507**, 214 (2001); E. Ma, Phys. Rev. D **66**, 117301 (2002); W. Grimus and L. Lavoura, Phys. Lett. B **572**, 189 (2003); see also P. F. Harrison and W. G. Scott, Phys. Lett. B **547**, 219 (2002); S. Choubey and W. Rodejohann, Eur. Phys. J. C **40**, 259 (2005); I. Aizawa, T. Kitabayashi and M. Yasue, Phys. Rev. D **72**, 055014 (2005).
 - [6] T. Schwetz, Acta Phys. Polon. B **36**, 3203 (2005), hep-ph/0510331.
 - [7] W. Grimus *et al.*, Nucl. Phys. B **713**, 151 (2005); R. N. Mohapatra, JHEP **0410**, 027 (2004); K. Matsuda and H. Nishiura, Phys. Rev. D **72**, 033011 (2005); R. N. Mohapatra and W. Rodejohann, Phys. Rev. D **72**, 053001 (2005); Y. H. Ahn *et al.*, hep-ph/0602160.
 - [8] A. S. Joshipura, hep-ph/0512252.
 - [9] F. Vissani, hep-ph/9708483; V. D. Barger, S. Pakvasa, T. J. Weiler and K. Whisnant, Phys. Lett. B **437**, 107 (1998); A. J. Baltz, A. S. Goldhaber and M. Goldhaber, Phys. Rev. Lett. **81**, 5730 (1998); H. Georgi and S. L. Glashow, Phys. Rev. D **61**, 097301 (2000); I. Stancu and D. V. Ahluwalia, Phys. Lett. B **460**, 431 (1999).
 - [10] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530**, 167 (2002).

- [11] For an overview, see S. Hannestad, Nucl. Phys. Proc. Suppl. **145**, 313 (2005), hep-ph/0412181.
- [12] S. Choubey, hep-ph/0509217.
- [13] Note that the form of the mass matrix Eq. (13) is gener-

ated at a scale of order TeV. Hence, the running of the mass and mixing parameters to low scale m_Z is rather moderate, in particular for small neutrino masses.